

NO EVIDENCE FOR CLASSICAL CEPHEIDS AND A NEW DWARF GALAXY
BEHIND THE GALACTIC DISK¹P. PIETRUKOWICZ², A. UDALSKI², M. K. SZYMAŃSKI², I. SOSZYŃSKI², G. PIETRZYŃSKI^{2,3}, L. WYRZYKOWSKI²,
R. POLESKI^{2,4}, K. ULACZYK^{2,5}, J. SKOWRON², P. MRÓZ², M. PAWLAK², AND S. KOZŁOWSKI²*Draft version November 30, 2015*

ABSTRACT

Based on data from the ongoing OGLE Galaxy Variability Survey (OGLE GVS) we have verified observed properties of stars detected by the near-infrared VVV survey in a direction near the Galactic plane at longitude $l \approx -27^\circ$ and recently tentatively classified as classical Cepheids belonging to a, hence claimed, dwarf galaxy at a distance of about 90 kpc from the Galactic Center. Three of four stars are detected in the OGLE GVS I -band images. We show that two of the objects are not variable at all and the third one with a period of 5.695 d and a nearly sinusoidal light curve of an amplitude of 0.5 mag cannot be a classical Cepheid and is very likely a spotted object. These results together with a very unusual shape of the K_s -band light curve of the fourth star indicate that very likely none of them is a Cepheid and, thus, there is no evidence for a background dwarf galaxy. Our observations show that a great care must be taken when classifying objects by their low-amplitude close-to-sinusoidal near-infrared light curves, especially with a small number of measurements. We also provide a sample of high-amplitude spotted stars with periods of a few days that can mimic pulsations and even eclipses.

Subject headings: galaxies: dwarf – Galaxy: disk – stars: variables: Cepheids

1. THE TARGET OBJECTS

Recently Chakrabarti et al. (2015) reported on the detection of four classical Cepheids clustered in angle (within one degree) and distance (on average 90 kpc from the Galactic Center) in a direction near the Galactic plane at longitude $l \approx -27^\circ$. They used near-infrared data from the ESO public survey VISTA Variables in the Via Lactea (VVV, Minniti et al. 2010). Their search for periodic variables was based on time-series photometry in the K_s band with only 32 epochs per star across the VVV disk area. For each of the star classified by them as a Cepheid they provide single epoch VVV photometry in the JHK_s bands and extinction values derived from the color excess assuming the Cardelli et al. (1989) extinction law. Finally, they calculate distances to the objects using a formula given in Feast et al. (2014). The obtained mean distance of 90 kpc and clumped location in the sky led them to a conclusion that the stars are associated with a previously unknown dwarf galaxy. In our work, we show that to conclude on the presence of such a galaxy one has to be absolutely sure that the observed stars are of particular type and are indeed located at a similar distance behind the Galactic disk.

2. ANALYSIS OF THE OGLE DATA

The OGLE project in its fourth phase of operation (OGLE-IV, Udalski et al. 2015) conducts, among oth-

ers, a survey of the Galactic disk area visible from Las Campanas Observatory, Chile—the OGLE Galaxy Variability Survey (OGLE GVS). It covers 2/3 of the Galactic plane with longitudes from -170° to $+60^\circ$ and latitudes $-3^\circ \lesssim b \lesssim +3^\circ$. Regular monitoring with a cadence of 1–2 days, called the shallow OGLE GVS, started in 2013. It spans a magnitude range $10 < I < 19$. A complementary, deeper survey started in 2010 and maps the Galactic plane area down to $I \approx 22$ mag. The data will be extremely useful for studying the Milky Way structure and properties of many variable objects (cf. Pietrukowicz et al. 2013a,b). However, at the moment of writing (June 2015) sufficient amount of data have been collected to find information on selected Galactic plane objects, such as the candidate Cepheids reported in Chakrabarti et al. (2015).

For clarity, it is very important to note that in their paper, Chakrabarti et al. (2015) made the following errors in IDs, coordinates, and finding charts for reported objects, later corrected in private communication with us: (1) everywhere in their paper the correct ID for the second reported star should be VVV J162231.35-512346.9 (not VVV J162328.18-513230.4) and thus the Galactic coordinates given in their Table 1 are also incorrect; (2) the finding chart for the second object is wrong, while the K_s -band light curve shown in their Fig. 1 is correct; (3) in the same figure, they labeled the fourth object using the ID of the third one. Our Table 1 gives correct positions of the investigated objects. Among minor things presented in Chakrabarti et al. (2015) in a different than common manner are the finding charts in Galactic coordinates instead of equatorial coordinates and an opposite direction of increasing Galactic longitude in their Fig. 1.

Once we made sure of the positions of the reported objects, we found three of the four stars, hereafter denoted as S1, S2, S3, and S4, in deep OGLE GVS I -band frames (see Fig. 1). Only object S4 located about $0^\circ 77$

¹ Based on observations obtained with the 1.3-m Warsaw telescope at the Las Campanas Observatory of the Carnegie Institution for Science

² Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland

³ Universidad de Concepción, Departamento de Astronomía, Casilla 160-C, Concepción, Chile

⁴ Department of Astronomy, Ohio State University, 140 W. 18th Ave., Columbus, OH 43210, USA

⁵ Department of Physics, University of Warwick, Coventry CV4 7AL, UK

TABLE 1
DATA ON INDIVIDUAL TARGET STARS

Star	VVV ID	l [$^{\circ}$]	b [$^{\circ}$]	OGLE field	OGLE ID	P_{OGLE} [d]
S1	J162559.36-522234.0	-27.597	-2.237	GD1126.25	6394	no
S2	J162231.35-512346.9	-27.273	-1.168	GD1133.28	15388	no
S3	J162119.39-520233.3	-27.862	-1.494	GD1133.13	8832	5.6950(3) or $\times 2$
S4	J161542.47-494439.0	-26.888	+0.768	GD1139.19	-	-




FIG. 1.— Images showing the four target stars (in the centers of the circles) in four different bands: K_sJZ from the 4.1-m VISTA telescope and in I from the 1.3-m OGLE telescope. Each image is $30'' \times 30''$. North is up and East to the left. Only the last target is not visible in the OGLE GVS data.

from the Galactic plane in a highly reddened area is not visible. Star S1 is bright enough to be verified for variability in the shallow OGLE GVS images, but stars S2 and S3 required additional deep observations taken in May–June 2015. The number of I -band measurements for S1, S2, and S3 is 61, 48, and 44, respectively. The photometry was extracted using the Difference Image Analysis (DIA) method with an image subtraction algorithm implemented by Woźniak (2000) to the OGLE images of dense stellar fields. Udalski et al. (2015) gives full details on the data reduction techniques. In Fig. 2, we show light curves of the three detected stars, while in Fig. 3, we present power spectra obtained with the




FIG. 2.— Time-domain (left panels) and phased (right panels) I -band light curves of the three stars detected in the OGLE Galaxy Variability Survey. Objects S1 and S2 clearly do not phase with the periods given in Chakrabarti et al. (2015). Deep OGLE observations confirm light variations in star S3 with a period of 5.695 d (or twice longer), but their shape is different than expected for fundamental-mode classical Cepheids. Observations from 2011 for this object are marked in red to show a likely amplitude change.

ANOVA statistics (Schwarzenberg-Czerny 1996). Stars S1 and S2 look to be constant. There is no significant period in their power spectra. Light curves clearly do not phase with the periods given in Chakrabarti et al. (2015). If the two objects were indeed Cepheids with pulsation periods of 3–4 d as proposed on the base of the K_s -band observations, we would see them in the I band as large-amplitude variables ($\Delta I \gtrsim 0.2$ mag). Objects S1 and S2 either are constant stars or have amplitudes of at most a few hundredths of magnitude.

Deep OGLE GVS observations confirm the variable nature of object S3 with a period around 5.69 d noted in Chakrabarti et al. (2015). The power spectrum based on the optical data shows a very strong signal at $P = 5.695$ d and also $2P = 11.390$ d. The shape of the light curve is close to a sinusoid, but its stability is under question. It is likely that the I -band amplitude changed from about 0.35 mag in 2011 to 0.50 mag in 2015. The value for 2011




FIG. 3.— ANOVA power spectra of the three objects observed by OGLE GVS. No periodic signal is detected in stars S1 and S2. Object S3 shows strong peaks at frequencies $1/5.695$ c/d and $1/11.39$ c/d.




FIG. 4.— Fourier series with increasing order from $n = 1$ to $n = 3$ fitted to the I -band light curve of star S3 folded with the period $P = 5.695$ d. Full amplitudes A are given to each fit. The fit with $n = 2$ seems to be optimal.

is, however, uncertain, since only eight images were taken that year. In Fig. 4, we show that the optimal Fourier fit to the I -band data is of an maximal order of $n = 2$, assuming $P = 5.695$ d and stability of the light curve. For $n = 2$ we obtain the following Fourier parameters: $A_0 = 19.593 \pm 0.009$, $A_1 = 0.251 \pm 0.016$, $A_2 = 0.055 \pm 0.016$, $\phi_1 = 5.03 \pm 0.06$, $\phi_2 = 3.49 \pm 0.29$. From this we find the following combinations: $R_{21} = A_2/A_1 = 0.219 \pm 0.065$




FIG. 5.— Fourier combination ϕ_{21} vs. $\log P$ for fundamental-mode (left panels) and first-overtone Cepheids (right panels) from the LMC (upper panels) and SMC (lower panels). Star S3 with a nearly sinusoidal light curve cannot be a fundamental-mode pulsator.




FIG. 6.— Phase difference between minimum and maximum brightness for fundamental-mode classical Cepheids with I -band amplitudes > 0.4 mag from SMC (left panel) and LMC (right panel). The observed abrupt phase difference at about 10 d is related to the effect of Hertzsprung progression. Star S3 with $P = 5.695$ d and nearly sinusoidal 0.5-mag light curve is a clear outlier.

and $\phi_{21} = \phi_2 - 2\phi_1 = 5.99 \pm 0.31$. In the case of $n = 3$, we would obtain: $A_0 = 19.591 \pm 0.009$, $A_1 = 0.249 \pm 0.016$, $A_2 = 0.053 \pm 0.016$, $A_3 = 0.043 \pm 0.016$, $\phi_1 = 5.03 \pm 0.06$, $\phi_2 = 3.59 \pm 0.31$, $\phi_3 = 5.09 \pm 0.36$, and hence: $R_{21} = A_2/A_1 = 0.213 \pm 0.066$, $\phi_{21} = \phi_2 - 2\phi_1 = 6.09 \pm 0.33$, $R_{31} = A_3/A_1 = 0.173 \pm 0.065$, $\phi_{31} = \phi_3 - 3\phi_1 = 2.56 \pm 0.41$. All above uncertainties were calculated from the standard propagation formula for independent variables. The combination ϕ_{21} is the best parameter to distinguish between the fundamental-mode and first-overtone pulsators with periods around 5.7 d ($\log P \approx 0.76$). The value of ϕ_{21} , here very similar for $n = 2$ and $n = 3$, is evidently too high for fundamental-mode Cepheids at this period (see Fig. 5), which light curves have characteristic saw-tooth shapes (Soszyński et al. 2008, 2010). To support our claim, in Fig. 6, we present how fast high-amplitude classical Cepheids increase their brightness from minimum to maximum. For stars with periods




FIG. 7.— Period-amplitude diagram for known first-overtone Cepheids from three different environments (from top to bottom): of our Galaxy (ASAS data, Pojmański 2002; Sitek & Pojmański 2014), LMC (OGLE data, Soszyński et al. 2008), and SMC (OGLE data, Soszyński et al. 2010). Star S3 with the period $P = 5.695$ d has evidently too high amplitude to be such a type of pulsator.




FIG. 9.— Fourier combinations R_{21} , R_{31} , ϕ_{21} , and ϕ_{31} in function of period for star S3 (red points) in comparison to the three spotted stars presented in Fig. 8. Black points represent separate seasons of observations. The parameters describing the light curve of S3 are in agreement with spotted objects.




FIG. 8.— Examples of I -band light curves of three spotted stars from OGLE. Identification names, equatorial coordinates, and periods are given for each object. Two selected seasons are shown with different colors. Light curves of spotted stars may have amplitudes exceeding 0.5 mag and may mimic pulsating stars and even eclipsing binaries if observed for a short period of time.

between 2 and 7 d it takes about 20% of the period, in contrast to 40% in S3.

The close-to-sinusoidal light curve shape of S3, or the ϕ_{21} value of 6.0 ± 0.3 , could indicate a first-overtone Cepheid, but I -band amplitudes as high as 0.5 mag

are not observed at long periods in this type of pulsators (see Fig. 7). Moreover, longest observed periods in first-overtone Cepheids tend to shorten with metallicity. For instance, periods in this type of Cepheids from the Large Magellanic Cloud are not longer than 5.9 d (Soszyński et al. 2008), while from more metal-poor Small Magellanic Cloud are not even longer than 4.5 d (Soszyński et al. 2010). In none of known environments, I -band amplitudes of the first-overtone Cepheids with periods > 2 d exceed 0.35 mag. A dwarf galaxy is expected to host old and metal-poor stars, such as RR Lyrae type variables and anomalous Cepheids rather than classical ones.

On the other hand, amplitudes of 0.5 mag or higher are observed in numerous chromospherically active stars, such as of RS CVn type. In Fig. 8, we show three examples of spotted stars with periods of a few days found in the OGLE database. Light curves of such stars change slowly over years and very often may mimic pulsations and eclipses if observed for a short period of time. Based on the observed properties of the light curve, we conclude that object S3 is very likely a spotted star with the period of 5.695 d, but the period of 11.390 d cannot be excluded. See comparison of the Fourier combinations for S3 and spotted stars in Fig. 9.

Object S4, which is too faint to be detected in the deep OGLE GVS images, according to Chakrabarti et al. (2015) has a period of 13.9 d and a skewed K_s -band light curve with the rising part about twice longer than the fading one. Such a shape is not observed in classical Cepheids. Soszyński et al. (2005) normalized near-IR photometry for 30 Galactic and 31 LMC fundamental-mode Cepheids with periods mostly longer than 10 d and showed that their K -band light curves are very homogeneous and nearly symmetric. Pejcha & Kochanek (2012)

performed a global model for 287 classical Cepheids with $P > 10$ d based on thousands of multiband observations. According to their work K -band light curves of stars with periods of 13–14 d have the rise and fall time scales equal. K -band light curves of long-period classical Cepheids presented in Persson et al. (2004), Monson & Pierce (2011), and Macri et al. (2015) are practically symmetric and none of them shows the rising branch twice longer than the fading one. Some of the presented light curves have a small number of points in K and their scatter is comparable with the amplitude, like OGLE-LMC-CEP-2562 in Macri et al. (2015) and S4 here.

3. CONCLUSIONS

Our observations show that none of the three VISTA objects detected in the OGLE GVS data is a classical Cepheid. Objects S1 and S2 do not look to be variable at all. Object S3 is indeed variable, but it has a nearly sinusoidal light curve with a high amplitude of 0.5 mag

in the I band, likely not stable in time. This is not observed in classical Cepheids in any environment. S3 is very likely a spotted star of RS CVn type. Although we are not able to detect star S4, its K_s -band light curve with $P = 13.9$ d and the twice longer rising part than the fading one would be very unusual for a fundamental-mode classical Cepheid. Based on the above findings we conclude that there is no evidence for the presence of a dwarf galaxy behind the Galactic disk in the observed direction. Our results show that a great care must be taken when classifying variable objects based on low-amplitude near-infrared light curves, especially when composed of a small number of data points.

The OGLE project has received funding from the National Science Centre, Poland, grant MAESTRO 2014/14/A/ST9/00121 to AU. This work has been also supported by the Polish Ministry of Sciences and Higher Education grants No. IP2012 005672 under the Iuventus Plus program to PP and No. IdP2012 000162 under the Ideas Plus program to IS.

REFERENCES

- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Chakrabarti, S., Saito, R., Quillen, A., et al. 2015, *ApJ*, 802, L4
- Feast, M. W., Menzies, J. W., Matsunaga, N., & Whitelock, P. A. 2014, *Nature*, 509, 342
- Macri, L. M., Ngeow, C.-C., Kanbur, S. M., Mahzooni, S., & Smitka, M. T. 2015, *AJ*, 149, 117
- Monson, A. J., & Pierce, M. J. 2011, *ApJS*, 193, 12
- Minniti D., Lucas, P. W., Emerson, J. et al. 2010, *New Astronomy*, 15, 433
- Pejcha, O., & Kochanek, C. S. 2012, *ApJ*, 748, 107
- Persson, S. E., Madore, B. F., Krzemiński, W., Freedman, W. L., Roth, M., & Murphy, D. C. 2004, *AJ*, 128, 2239
- Pietrukowicz, P., Mróz, P., Soszyński, et al. 2013a, *Acta Astron.*, 63, 115
- Pietrukowicz, P., Dziembowski, W. A., Mróz, P., et al. 2013b, *Acta Astron.*, 63, 379
- Pojmański, G. 2002, *Acta Astron.*, 52, 397
- Schwarzenberg-Czerny, A. 1996, *ApJ*, 460, L107
- Sitek, M., & Pojmański, G. 2014, *Acta Astron.*, 64, 115
- Soszyński, I., Gieren, W., & Pietrzyński, G. 2005, *PASP*, 117, 823
- Soszyński, I., Poleski, R., Udalski, A., et al. 2008, *Acta Astron.*, 58, 163
- Soszyński, I., Poleski, R., Udalski, A., et al. 2010, *Acta Astron.*, 60, 17
- Udalski, A., Szymański, M., & Szymański, G. 2015, *Acta Astron.*, 65, 1
- Woźniak, P. R. 2000, *Acta Astron.*, 50, 421